arXiv:1108.4643v1 [hep-th] 23 Aug 2011

Coulomb Gas and Sine-Gordon Model in Arbitrary Dimension

I. Nándori[1]

[1]*Institute of Nuclear Research, P.O.Box 51, H-4001 Debrecen, Hungary*

Exact phase structure of the neutral Coulomb gas has been mapped out by the functional renormalization group study of the equivalent sine-Gordon field theory in arbitrary dimension. The high temperature fixed point is obtained and it is shown that the Coulomb gas has a single phase for $d > 2$. Renormalization group equations of the 3-dimensional sine-Gordon model are compared to those of the vortex-loop gas i.e. the gas of topological defects of the 3-dimensional isotropic XY spin model which demonstrates that the two models belong to different universality classes.

PACS numbers: 11.10.Hi, 05.70.Fh, 64.60.-i, 05.10.Cc

I. INTRODUCTION

Universality which expresses the idea that different microscopic physics can give rise to the same scaling behavior at a phase transition, is one of the most powerful tool of critical phenomena. For example, if two different models can be mapped onto each other then they should belong to the same universality class hence by the study of one model one can obtain the critical behavior of the other one. If the critical behaviors of two models agree or disagree it is possible to decide whether they belong to the same universality class. Furthermore, models of the same class of universality can be used to test and compare various types of methods suitable for the study of critical behavior. A canonical example for model equivalences is the sine-Gordon (SG) scalar field theory defined by the Euclidean action [1–4]

$$S = \int d^d x \left[\frac{1}{2}(\partial_\mu \varphi_x)^2 + u\cos(\beta\varphi_x)\right], \tag{1}$$

which belongs to the universality class of the neutral Coulomb gas (CG) [5] and the isotropic XY classical spin model [1] for $d = 2$. They undergo a Kosterlitz-Thouless-Berezinski [6] (KTB) phase transition which was well reproduced in the framework of the O(N) symmetric scalar field theory for $d = 2$ and $N = 2$ [7]. Furthermore, the SG scalar theory is the bose form of the 2-dimensional fermionic Thirring model [8] and the bosonic counterpart of multi-flavor Quantum Electrodynamics and multi-color Quantum Chromodynamics are also SG type models [9]. Although bosonization is well established only in $d = 2$ dimensions, it is known that the mapping between the SG theory (1) and the neutral CG holds in arbitrary dimension [1, 3].

The goal of this work is (i) to consider possible equivalences between CG, SG, XY models in $d \neq 2$ dimensions, (ii) to map out the exact phase structure of the d-dimensional neutral CG by the functional renormalization group (RG) study of the equivalent SG scalar theory.

II. RG STUDY OF THE SG MODEL

By the inclusion of the wave function renormalization in the functional RG study of the SG scalar field theory performed in arbitrary dimension we go beyond the approximations used in previous works [2–4]. We use the effective average action functional RG method [10] where the evolution equation reads as

$$k\partial_k \Gamma_k[\varphi] = \frac{1}{2}\mathrm{Tr}[(R_k + \partial^2_\varphi \Gamma_k[\varphi])^{-1}\, k\partial_k R_k], \tag{2}$$

where k is the running momentum cutoff, the trace Tr stands for the integration over all momenta. We introduce the power-law regulator function $R_k = p^2\left(k^2/p^2\right)^b$ where $b \geq 1$. Let us note, that the field carries a dimension for $d \neq 2$ thus the frequency of the SG model (1) becomes a dimensionful parameter for $d \neq 2$, i.e. $\beta^2 = k^{2-d}\tilde\beta^2$ where $\tilde\beta$ is dimensionless. Eq. (2) has been solved over the functional subspace spanned by the ansatz for the SG model

$$\Gamma_k = \int d^d x \left[\frac{1}{2}\frac{1}{\tilde\beta_k^2}(\partial_\mu \varphi_x)^2 + u_k \cos(k^{\frac{2-d}{2}}\varphi_x)\right], \tag{3}$$

where the local potential contains a single Fourier mode and the field has been rescaled as $\varphi \to \varphi/\tilde\beta$. In case of the d-dimensional SG model it is convenient to introduce a dimensionless variable $\tilde\varphi = k^{(2-d)/2}\varphi$, then (3) reads as

$$\Gamma_k = \int d^d x \left[\frac{1}{2} z_k (\partial_\mu \tilde\varphi_x)^2 + V_k(\tilde\varphi_x)\right], \quad V_k(\tilde\varphi_x) = u_k \cos(\tilde\varphi_x) \tag{4}$$

where $z_k \equiv 1/\beta_k^2 = 1/(k^{2-d}\tilde\beta_k^2)$ stands for the field-independent wave-function renormalization which has a dimension of k^{d-2}. The RG equation has been modified if a dimensionless field variable $\tilde\varphi$ has been introduced,

$$k\partial_k \Gamma_k[\tilde\varphi] = \frac{1}{2}\mathrm{Tr}\left[\frac{k\partial_k R_k}{R_k + k^{d-2}\Gamma''_k[\tilde\varphi]}\right] \tag{5}$$

with the notation $' = \partial/\partial\tilde\varphi$. Let us note the term $-[(d-2)/2]\tilde\varphi\partial_{\tilde\varphi}$ is neglected on l.h.s. of (5) because it does not give contribution to the periodic part of the action [3] and the blocked potential $V_k(\tilde\varphi_x)$ contains only periodic interactions (and z_k is field-independent). Although RG transformations generate higher harmonics, we use the ansatz (4) which contains a single Fourier mode since in case of the 2-dimensional SG model [11] it was found to be an appropriate approximation [12]. Then Eq. (5) leads to the evolution equations for the blocked potential $V_k(\tilde\varphi)$ and the wave-function renormalization z_k,

$$\partial_k V_k = \frac{1}{2}\int \frac{d^dp}{(2\pi)^d}\mathcal{D}_k\, k\partial_k R_k, \tag{6}$$

$$k\partial_k z_k = \left(\frac{1}{2\pi}\int_0^{2\pi} d\tilde\varphi\right)(k^{2-d}V_k''')^2 \int \frac{d^dp}{(2\pi)^d} \mathcal{D}_k^2\, k\partial_k R_k \left(\frac{2}{d}\frac{\partial^2\mathcal{D}_k}{\partial p^2 \partial p^2}p^2 + \frac{\partial \mathcal{D}_k}{\partial p^2}\right), \tag{7}$$

with $\mathcal{D}_k = (z_k k^{2-d}p^2 + R_k + k^{2-d}V_k'')^{-1}$. Let us note since the l.h.s of (7) is independent of the field, a projection onto the field-independent subspace has been introduced on the r.h.s of (7). The scale k covers the momentum interval from the high-energy/ultraviolet (UV) cutoff Λ to zero. Inserting the ansatz (4) into Eqs. (6), (7), flow equations for the dimensionful couplings are

$$k\partial_k u_k = \int_p \frac{k\partial_k R_k}{k^{2-d}u_k}\left(\frac{P-\sqrt{P^2-(k^{2-d}u_k)^2}}{\sqrt{P^2-(k^{2-d}u_k)^2}}\right), \tag{8}$$

$$k\partial_k z_k = \int_p \frac{k\partial_k R_k}{2}\Bigg[\frac{-(k^{2-d}u_k)^2 P(\partial_{p^2}P + \frac{2}{d}p^2\partial^2_{p^2}P)}{[P^2-(k^{2-d}u_k)^2]^{5/2}} + \frac{(k^{2-d}u_k)^2 p^2(\partial_{p^2}P)^2(4P^2+(k^{2-d}u_k)^2)}{d\,[P^2-(k^{2-d}u_k)^2]^{7/2}}\Bigg], \tag{9}$$

where $P = z_k k^{2-d}p^2 + R_k$ and $\int_p = \int dp\, p^{d-1}\Omega_d/(2\pi)^d$ with the d-dimensional solid angle Ω_d. Momentum integrals have to be performed numerically, except the linearized form of Eqs. (8), (9) around the Gaussian fixed point where analytical results available. The linearized RG with dimensionless couplings $\tilde u_k = k^{-d}u_k$, $\tilde z_k = k^{2-d}z_k$ reads as for $d = 1$ dimensions

$$(1+k\partial_k)\tilde u_k = \frac{1}{4b\sin(\frac{\pi}{2b})}\tilde z_k^{\frac{1-2b}{2b}}\tilde u_k + \mathcal{O}(\tilde u_k^2) \tag{10}$$

$$(-1+k\partial_k)\tilde z_k = -\frac{c_1(b)}{4\pi}\tilde z_k^{\frac{5-4b}{2b}}\tilde u_k^2 + \mathcal{O}(\tilde u_k^3) \tag{11}$$

with $c_1(b) = \frac{5\pi(55+4\pi(5b-18))}{48b\sin[5\pi/(2b)]}$. Eqs. (10), (11) have a non-trivial fixed point at $\tilde z_\star = [4b\sin(\pi/(2b)]^{2b/(1-2b)}$, $\tilde u_\star = (4\pi/c_1)\tilde z_\star^{3-5/(2b)}$. For $d = 2$, linearized RG equations are

$$(2+k\partial_k)\tilde u_k = \frac{1}{4\pi\tilde z_k}\tilde u_k + \mathcal{O}(\tilde u_k^2) \tag{12}$$

$$k\partial_k\tilde z_k = -\frac{c_2(b)}{8\pi}\tilde z_k^{\frac{2-2b}{b}}\tilde u_k^2 + \mathcal{O}(\tilde u_k^3) \tag{13}$$

with $c_2(b) = \frac{2\pi(b-2)(b-1)}{3b^2\sin[2\pi/b]}$ and result in a KTB type (i.e. infinite order) phase transition with $\tilde z_\star = 1/(8\pi)$ [11, 12]. Finally, for $d = 3$, linearized RG equations are

$$(3+k\partial_k)\tilde u_k = \frac{1}{8\pi b\sin(\frac{\pi}{2b})}\tilde z_k^{\frac{-1-2b}{2b}}\tilde u_k + \mathcal{O}(\tilde u_k^2) \tag{14}$$

$$(1+k\partial_k)\tilde z_k = -\frac{c_3(b)}{8\pi^2}\tilde z_k^{\frac{3-4b}{b}}\tilde u_k^2 + \mathcal{O}(\tilde u_k^3) \tag{15}$$

with $c_3(b) = \frac{\pi(3+4b(b-2))}{16b^2\sin[3\pi/(2b)]}$. Due to the tree-level scaling of $\tilde z_k$, the non-trivial fixed point appears for $d < 2$ and disappears for $d > 2$ in the RG flow. However, a "turning point" can be identified for $d > 2$ where the irrelevant coupling $\tilde u_k$ turns to a relevant one. For $d = 3$ the turning point is at $\tilde z_\star = [24\pi b\sin(\pi/(2b)]^{-2b/(1+2b)}$.

In general, RG equations (8) and (9) (exact for a single Fourier mode) have to be solved numerically. Important feature of the exact RG flow the emergence of a new low-energy/infrared (IR) fixed point related to the degeneracy of the blocked action. Namely, Eqs. (8), (9) become singular at the momentum scale where $\bar k - k^{2-d}u_k = 0$ with $\bar k = \min_{p^2} P = bk^2[\tilde z_k/(b-1)]^{1-1/b}$. Therefore, it is convenient to redefine the dimensionless coupling constant as $\bar u_k \equiv k^{2-d}u_k/\bar k = k^2\tilde u/\bar k$ which tends to one in case of degeneracy. Exact RG equations (8), (9) were solved for $b = 2$ and flow diagrams are plotted in Fig. 1 for $d = 1$, and in Fig. 2 for $d = 3$ dimensions. The phase structure is independent of b but we use $b = 2$ here since it was found to be an optimal renormalization scheme for SG type models [13]. The IR fixed point corresponds to the degeneracy was found in any dimensions. However, the $d = 1$ case possibly needs further improvement since its phase structure shows similarities to the QM anharmonic oscillator where spontaneous symmetry breaking is not allowed [14]. There the higher order terms in the gradient expansion handle the problem.



FIG. 1: Phase diagram of the SG and the equivalent CG models for $d = 1$ dimensions (similar RG flow can be drawn for $d < 2$). Arrows indicate the direction of the flow. Parameters $\bar u$, $\tilde z$ are related to the fugacity y and the temperature $\tilde T$.

Let us emphasize that only the exact RG flow with the inclusion of the wave function renormalization is suitable for the determination of the IR fixed point since the



FIG. 2: Phase diagram of the SG and the equivalent CG models for $d = 3$ dimensions which indicates a single phase for $d > 2$.

perturbative (truncated) RG equations are non-singular [2]. Thus any previous attempts based on perturbation theory [2] or by the usage of the local potential approximation [3, 4] were unable to determine the IR behavior in a reliable manner. Furthermore, the exact RG flow changes the position of the non-trivial fixed point obtained for $d = 1$. Indeed, for $b = 2$, linearized RG equations (10), (11) result in $1/\tilde{z}_\star = 2^{10/3} \approx 10.08$ and $\bar{u}_\star = \tilde{u}_\star/(2\sqrt{\tilde{z}_\star}) = 2/\sqrt{15} \approx 0.52$ which have been modified by the exact RG, see Fig. 1. Similarly, the turning point of the $d = 3$ case determined by the exact RG coincides with that of the linearized RG which is $1/\tilde{z}_\star = 4(3\pi)^{2/3} \approx 41.9$ only for small $\bar{u}$, see Fig. 2. Finally, let us mention that the comparison of exact and perturbative (linearized) RG flow can be used for optimizing the renormalization schemes. For $d = 1$ and $b = 1$ the non-trivial fixed point determined by the linearized RG ($1/\tilde{z}_\star = 16$ and $\bar{u}_\star = \tilde{u}_\star = 4/\sqrt{5} \approx 1.79$) is out of the range of validity of the exact flow ($\bar{u}_k = \tilde{u}_k \le 1$), hence it represents a drawback on the applicability of the RG scheme $b = 1$ for $d \neq 2$ dimensions.

III. PHASE STRUCTURE OF THE NEUTRAL CG

The CG model has been the subject of an intense study in last decades [2, 3, 5, 15–17] and there is a continuos interest in the usage of the SG representation of CG systems [2, 3, 5, 17] since it has an important relevance in various models of critical phenomena (e.g. superfluid transition in He films, melting of crystals in $d = 2$, arrays of Josephson junctions, roughening transition). However, the complete understanding of the critical behavior of the d-dimensional neutral CG which is a plasma of equal number of positive and negative charges is still lacking. For example, previous investigations (even for the well studied 2-dimensional CG model) were restricted to the case of low densities (limit of low fugacity) and there are different competing views of what happens to the neutral plasma at high densities [16, 17]. Thus, an RG study of the d-dimensional CG beyond the dilute gas approximation is required which is the goal of the present work.

Since the mapping between the CG and SG models holds in arbitrary dimension [3, 5] (and it is exact in case of point-like charges) the RG study of the d-dimensional SG model can be directly used to map out the phase structure of the CG model. In the framework of perturbation theory (limit of low fugacity) RG equations of the CG model in arbitrary dimension [15] reads as

$$\frac{dx}{dl} = -x(xy^2 + d - 2), \quad \frac{dy}{dl} = -y(x - d) \qquad (16)$$

with $\partial_l = -k\partial_k$, $y \sim \tilde{u}_k$ and $x \sim 1/\tilde{T} \sim 1/\tilde{z}_k$ where $\tilde{T}$ is the temperature and y is the fugacity. Using these identities Eq. (16) can be rewritten as

$$((d-2)+k\partial_k)\tilde{z}_k = -c_z\tilde{u}_k^2, \quad (d+k\partial_k)\tilde{u}_k = c_u\frac{1}{\tilde{z}_k}\tilde{u}_k, \quad (17)$$

with constants c_u, c_z and they are found to be similar to the linearized RG equations obtained for the SG model, e.g. the non-trivial fixed point of the 1-dimensional SG model can be identified in the flow generated by (17). In the sharp cutoff limit ($b \to \infty$) the linearized RG equation derived for the Fourier amplitude of the SG model reduces to $(d+k\partial_k)\tilde{u}_k = \Omega_d\tilde{u}_k/(\tilde{z}_k 2(2\pi)^d)$ which is identical to the second equation of (17). However, the sharp cutoff does not support to the derivative expansion at higher order, i.e. for $b \to \infty$ the multiplicative constant $c_d(b)$ of the RG equation obtained for $\tilde{z}_k$ becomes infinity.

Most important new results of the exact RG flow are the existence of the high temperature ($\bar{u}_\star = 1$, $1/\tilde{z}_\star = 0$) and the absence of new further non-trivial fixed points. Let us note that, the perturbative (truncated) RG has a non-singular structure hence it is not able to recover the degenerate potential therefore it is not suitable to observe the high temperature fixed point [15]. Moreover, the exact RG changes the position of the non-trivial fixed point for $d < 2$ and the turning point for $d > 2$. Furthermore, since the mapping between the SG and CG models is exact for point-like charges the exact RG flow indicates a single phase for the CG for $d > 2$.

IV. ISOTROPIC CLASSICAL XY SPIN MODEL

The partition function of the d-dimensional isotropic classical XY spin model [1] can be expressed in terms of topological excitations of the original degrees of freedom. For $d = 2$ dimensions, the dual theory is the vortex gas which is known to belong to the class of universality of the neutral CG [18, 19, 22]. Two-dimensional generalized models are well known where both the CG and the vortex gas are included as particular limiting cases [18, 19, 22] and are self-dual under the duality transformation. For

$d=3$, the dual theory is the gas of interacting vortex loops [20, 21] (i.e. the lattice CG [22]). Corresponding flow equations have been derived for the parameters K (i.e. the coupling between the spins) and y (i.e. the fugacity of the vortex loops) by real-space RG method ($a \sim 1/k$) in the limit of low fugacity [3, 20, 21],

$$a\partial_a K = K - \frac{4\pi^3}{3}K^2 y, \qquad a\partial_a y = \left(6 - \pi^2 K L\right) y, \quad (18)$$

where L approaches a constant in the IR limit ($L \to 1$) [20], or it is weakly divergent ($L \to \ln(a/a_c)+1$) [21]. Since $a\partial_a = -k\partial_k$ and by using the identities $y \equiv \tilde{u}$, $K \equiv 1/\tilde{z}$, RG equations of (18) are rewritten as

$$k\partial_k \tilde{z}_k = \tilde{z}_k - \frac{4\pi^3}{3}\tilde{u}_k^2, \quad k\partial_k \tilde{u}_k = \left(\frac{1}{\tilde{z}_k}\frac{\pi^2 L}{2} - 3\right)\tilde{u}_k, \quad (19)$$

which can be compared to the linearized RG obtained for the SG (14), (15) and the CG (17) models in $d=3$ dimensions. The only disagreement is the sign of the tree-level scaling term of $\tilde{z}_k$ but it has important consequences since the RG flow of the vortex-loop gas has a non-trivial fixed point which is absent in case of the 3-dimensional SG and CG models. Therefore, it demonstrates that the vortex-loop gas has a different critical behavior, thus it belongs to a class of universality differs from that of the SG and CG models for $d \neq 2$ (for $d=2$, couplings $K, \tilde{z}$ have no tree-level scaling thus SG, CG, XY models are in the same universality class). Let us emphasize that RG equations are compared at the linearized level, however the exact RG study of the SG and CG models is required since it shows the absence of new non-trivial fixed points, thus there is no room to find a mapping between the parameters of the vortex-loop gas and the 3-dimensional CG which could produce the same phase structure.

V. SUMMARY

In this work, the exact phase structure of the neutral CG was mapped out by the functional RG study of the equivalent SG scalar field theory in arbitrary dimension. By the inclusion of the wave function renormalization and the determination of the exact RG flow for a single Fourier mode, we went beyond the approximations used previously. The high temperature fixed point of the d-dimensional CG was determined and shown that the perturbative RG is not suitable to recover it. The position of the non-trivial fixed point of the RG flow for $d<2$ and the turning point for $d>2$ dimensions given by the exact flow is compared to the linearized one. The non-trivial fixed point of the CG for $d<2$ shows similarities to the Wilson-Fisher fixed point of the $O(N)$ model. The absence of new further non-trivial fixed points were also demonstrated which results in a single phase for the neutral CG with point-like charges for $d>2$. This has two consequences, (i) no further phase transition can be identified at high densities for the neutral CG as opposed to e.g. [16, 17], (ii) there is no way to find a mapping between the phase structure of the neutral 3-dimensional CG and the vortex-loop gas which is the gas of topological defects of the isotropic XY spin model for $d=3$, thus, it was demonstrated they belong to different universality classes.

Acknowledgement

This research was supported by the TÁMOP 4.2.1./B-09/1/KONV-2010-0007 project.

---

[1] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, Oxford, Clarendon (1989).

[2] G. Busiello, L. De Cesare, I. Rabuffo, Phys. Rev. B **32**, 5918 (1985); A. Caramico D'Auria, *et al.*, Physica A **274**, 410 (1999).

[3] I. Nándori, K. Sailer, U. D. Jentschura, G. Soff, Phys. Rev. D **69**, 025004 (2004);

[4] V. Pangon, arXiv:1008.0281 [hep-th].

[5] Stu Samuel, Phys. Rev. D **18**, 1916 (1978).

[6] V. L. Berezinskii, Zh. Eksp. Teor. Fiz. **61**, 1144 (1971) [Sov. Phys.-JETP **34**, 610 (1972); J. M. Kosterlitz, D. J. Thouless, J. Phys. C**6**, 1181 (1973).

[7] M. Gräter, C. Wetterich, Phys. Rev. Lett. **75**, 378 (1995); G. v. Gersdorff, C. Wetterich, Phys. Rev. B **64**, 054513 (2001).

[8] S. R. Coleman Phys. Rev. D **11**, 2088 (1975).

[9] S. Nagy, J. Polonyi, K. Sailer, Phys. Rev. D **70**, 105023 (2004); I. Nándori, J. Phys. A: Math. Gen. **39**, 8119 (2006); I. Nándori, Phys. Lett. B **662**, 302 (2008); S. Nagy, Phys. Rev. D **79**, 045004 (2009); J. Kovács, S. Nagy, I. Nándori, K. Sailer, JHEP **1101**, 126 (2011).

[10] C. Wetterich, Phys. Lett. B**301**, 90 (1993); T. R. Morris, Int. J. Mod. Phys. A **9**, 2411 (1994).

[11] I. Nándori, J. Polonyi, K. Sailer, Phys. Rev. D **63**, 045022 (2001); S. Nagy, K. Sailer, J. Polonyi, J. Phys. A **39**, 8105 (2006); S. Nagy, I. Nándori, J. Polonyi, K. Sailer, Phys. Lett. B **647**, 152 (2007); I. Nándori, S. Nagy, K. Sailer, A. Trombettoni, Phys. Rev. D **80**, 025008 (2009). V. Pangon, S. Nagy, J. Polonyi, K. Sailer, Phys. Lett. B **694**, 89 (2010); I. Nándori, S. Nagy, K. Sailer, A. Trombettoni, JHEP **1009**, 069 (2010).

[12] S. Nagy, I. Nándori, K. Sailer, J. Polonyi, Phys. Rev. Lett. **102**, 241603 (2009); S. Nagy, K. Sailer, arXiv:1012.3007 [hep-th].

[13] I. Nándori, arXiv:1008.2934.

[14] A.S. Kapoyannis, N. Tetradis, Phys. Lett. A**276**, 225 (2000); S. Nagy, K. Sailer, Annals Phys. **326**, 1839 (2011).

[15] J. M. Kosterlitz, J. Phys. C**10**, 3753 (1977).

[16] P. Minnhagen, Phys. Rev. Lett. **54**, 2351 (1985); P. Minnhagen, M. Wallin, Phys. Rev. B **40**, 5109 (1989); Y. Levin, X. Li, M.E. Fisher, Phys. Rev. Lett. **73**, 2716

(1994); S. Kragset, A. Sudbo, F.S. Nogueira, Phys. Rev. Lett. **92**, 186403 (2004); K. Borkje, S. Kragset, A. Sudbo, Phys. Rev. B **71**, 085112 (2005); S. Akhanjee, J. Rudnick, Phys. Rev. Lett. **105**, 047206 (2010);

[17] A. Diehl, M. C. Barbosa, Y. Levin, Phys. Rev. E **56**, 619 (1997); L. Samaj, B. Jancovici, J. Stat. Phys. **106**, 323 (2002); L. Samaj, J. Phys. A **36**, 5913 (2003); L, Mondaini, E.C. Marino, J. Stat. Phys. **118**, 767 (2005); L, Mondaini, E.C. Marino, J. Phys. A **39**, 967 (2006); G. Téllez, J. Stat. Phys. **126**, 281 (2007); L. Samaj, J. Stat. Phys. **128**, 569 (2007);

[18] B. Nienhuis in *Phase Transitions and Critical Phenomena*, Vol. 11 ed. by C. Domb, J.L. Lebowitz (Academic Press, London, 1987), 1-53.

[19] K. Huang, J. Polonyi, Int. J. Mod. Phys. **A6** 409 (1991).

[20] D. Nelson and D. Fisher, Phys. Rev. **B16** (1977) 4945.

[21] S. R. Shenoy, Phys. Rev. **B40** (1989) 5056.

[22] J.V. Jose, L.P. Kadanoff, S. Kirkpatrick, D.R. Nelson, Phys. Rev. B **16**, 1217 (1977); P. Minnhagen, B. J. Kim, H. Weber, Phys. Rev. Lett. **87**, 037002 (2001).